

18

# PERIODIC TABLE OF THE ELEMENTS

1

1	2	13	14	15	16	17	18
<b>H</b> 1.008	<b>He</b> 4.00	<b>B</b> 10.81	<b>C</b> 12.01	<b>N</b> 14.01	<b>O</b> 16.00	<b>F</b> 19.00	<b>Ne</b> 20.18
3	4	5	6	7	8	9	10
<b>Li</b> 6.94	<b>Be</b> 9.01	<b>B</b> 10.81	<b>C</b> 12.01	<b>N</b> 14.01	<b>O</b> 16.00	<b>F</b> 19.00	<b>Ne</b> 20.18
11	12	13	14	15	16	17	18
<b>Na</b> 22.99	<b>Mg</b> 24.30	<b>Al</b> 26.98	<b>Si</b> 28.09	<b>P</b> 30.97	<b>S</b> 32.06	<b>Cl</b> 35.45	<b>Ar</b> 39.95
19	20	21	22	23	24	25	26
<b>K</b> 39.10	<b>Ca</b> 40.08	<b>Sc</b> 44.96	<b>Ti</b> 47.87	<b>V</b> 50.94	<b>Cr</b> 52.00	<b>Mn</b> 54.94	<b>Fe</b> 55.85
37	38	39	40	41	42	43	44
<b>Rb</b> 85.47	<b>Sr</b> 87.62	<b>Y</b> 88.91	<b>Zr</b> 91.22	<b>Nb</b> 92.91	<b>Mo</b> 95.95	<b>Tc</b> 98.91	<b>Ru</b> 101.07
55	56	57-71	72	73	74	75	76
<b>Cs</b> 132.91	<b>Ba</b> 137.33	* Lanthanoids	<b>Hf</b> 178.49	<b>Ta</b> 180.95	<b>W</b> 183.84	<b>Re</b> 186.21	<b>Os</b> 190.23
87	88	89-103	104	105	106	107	108
<b>Fr</b> 87	<b>Ra</b> 88	† Actinoids	<b>Rf</b> 104	<b>Db</b> 105	<b>Sg</b> 106	<b>Bh</b> 107	<b>Hs</b> 108
111	112	113	114	115	116	117	118
<b>Cn</b> 112	<b>Cn</b> 112	<b>Nh</b> 113	<b>Fl</b> 114	<b>Mc</b> 115	<b>Lv</b> 116	<b>Ts</b> 117	<b>Og</b> 118
119	120	121	122	123	124	125	126
<b>Uu</b> 119	<b>Uu</b> 120	<b>Uu</b> 121	<b>Uu</b> 122	<b>Uu</b> 123	<b>Uu</b> 124	<b>Uu</b> 125	<b>Uu</b> 126

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
<b>La</b> 138.91	<b>Ce</b> 140.12	<b>Pr</b> 140.91	<b>Nd</b> 144.24	<b>Pm</b>	<b>Sm</b> 150.36	<b>Eu</b> 151.97	<b>Gd</b> 157.25	<b>Tb</b> 158.93	<b>Dy</b> 162.50	<b>Ho</b> 164.93	<b>Er</b> 167.26	<b>Tm</b> 168.93	<b>Yb</b> 173.05	<b>Lu</b> 174.97
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
<b>Ac</b> 232.04	<b>Th</b> 232.04	<b>Pa</b> 231.04	<b>U</b> 238.03	<b>Np</b>	<b>Pu</b>	<b>Am</b>	<b>Cm</b>	<b>Bk</b>	<b>Cf</b>	<b>Es</b>	<b>Fm</b>	<b>Md</b>	<b>No</b>	<b>Lr</b>

\*Lanthanoids

†Actinoids



## AP<sup>®</sup> CHEMISTRY EQUATIONS AND CONSTANTS

Throughout the exam the following symbols have the definitions specified unless otherwise noted.

L, mL = liter(s), milliliter(s)  
 g = gram(s)  
 nm = nanometer(s)  
 atm = atmosphere(s)

mm Hg = millimeters of mercury  
 J, kJ = joule(s), kilojoule(s)  
 V = volt(s)  
 mol = mole(s)

### ATOMIC STRUCTURE

$$E = h\nu$$

$$c = \lambda\nu$$

$E$  = energy  
 $\nu$  = frequency  
 $\lambda$  = wavelength

Planck's constant,  $h = 6.626 \times 10^{-34}$  J s

Speed of light,  $c = 2.998 \times 10^8$  m s<sup>-1</sup>

Avogadro's number =  $6.022 \times 10^{23}$  mol<sup>-1</sup>

Electron charge,  $e = -1.602 \times 10^{-19}$  coulomb

### EQUILIBRIUM

$$K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } a A + b B \rightleftharpoons c C + d D$$

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b}$$

$$K_a = \frac{[H^+][A^-]}{[HA]}$$

$$K_b = \frac{[OH^-][HB^+]}{[B]}$$

$$K_w = [H^+][OH^-] = 1.0 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log[H^+], \text{ pOH} = -\log[OH^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[A^-]}{[HA]}$$

$$\text{p}K_a = -\log K_a, \text{ p}K_b = -\log K_b$$

#### Equilibrium Constants

$K_c$  (molar concentrations)

$K_p$  (gas pressures)

$K_a$  (weak acid)

$K_b$  (weak base)

$K_w$  (water)

### KINETICS

$$[A]_t - [A]_0 = -kt$$

$$\ln[A]_t - \ln[A]_0 = -kt$$

$$\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$$

$$t_{1/2} = \frac{0.693}{k}$$

$k$  = rate constant

$t$  = time

$t_{1/2}$  = half-life

## GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$P_A = P_{\text{total}} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{\text{total}} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$D = \frac{m}{V}$$

$$KE_{\text{molecule}} = \frac{1}{2}mv^2$$

Molarity,  $M$  = moles of solute per liter of solution

$$A = \epsilon bc$$

$P$  = pressure

$V$  = volume

$T$  = temperature

$n$  = number of moles

$m$  = mass

$M$  = molar mass

$D$  = density

$KE$  = kinetic energy

$v$  = velocity

$A$  = absorbance

$\epsilon$  = molar absorptivity

$b$  = path length

$c$  = concentration

Gas constant,  $R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$

$= 0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1}$

$= 62.36 \text{ L torr mol}^{-1} \text{ K}^{-1}$

$1 \text{ atm} = 760 \text{ mm Hg} = 760 \text{ torr}$

STP =  $273.15 \text{ K}$  and  $1.0 \text{ atm}$

Ideal gas at STP =  $22.4 \text{ L mol}^{-1}$

## THERMODYNAMICS/ELECTROCHEMISTRY

$$q = mc\Delta T$$

$$\Delta S^\circ = \sum S^\circ_{\text{products}} - \sum S^\circ_{\text{reactants}}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ_{\text{products}} - \sum \Delta H_f^\circ_{\text{reactants}}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ_{\text{products}} - \sum \Delta G_f^\circ_{\text{reactants}}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K$$

$$= -nFE^\circ$$

$$I = \frac{q}{t}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \ln Q$$

$q$  = heat

$m$  = mass

$c$  = specific heat capacity

$T$  = temperature

$S^\circ$  = standard entropy

$H^\circ$  = standard enthalpy

$G^\circ$  = standard Gibbs free energy

$n$  = number of moles

$E^\circ$  = standard reduction potential

$I$  = current (amperes)

$q$  = charge (coulombs)

$t$  = time (seconds)

$Q$  = reaction quotient

Faraday's constant,  $F = 96,485$  coulombs per mole of electrons

$$1 \text{ volt} = \frac{1 \text{ joule}}{1 \text{ coulomb}}$$